



Development of a holistic strategy for the assessment of the quality of life in patients with breast cancer in the different stages of the disease [

Asociación de Psicooncología de Madrid,
2021-03-01

[info:eu-repo/semantics/article](#) [info:eu-repo/semantics/publishedVersion](#)
[Artículo revisado por pares](#)

Analítica

Introduction: Breast cancer is considered a chronic disease that has considerable impact on the quality of life (QoL) of the patients. Yet, there is not a totally satisfactory evaluation system reflecting the complexity of the breast cancer condition. Furthermore, most instruments are more addressed towards early-stage disease than advanced disease. **Objective:** To describe the methodology used to measure the QoL of life and coping strategies in a representative group of breast cancer patients including metastatic and non-metastatic patients. **Methodology:** Prospective study including patients with breast cancer at different stages of the disease recruited at the Medical Oncology Department of the Catalan Oncology Institute (ICO) previous informed consent. A protocol of approximately one hour face-to-face interview is run to collect sociodemographic information and to answer the questions of several QoL scales such as the QLCA-AFex Font, QLQ-C30, QLQ-BR23, HADS, DME, BRCS, MINI-MAC, LOT-R and OE questionnaires, complemented by a semi-structured interview. **Results:** From June 2017 to March 2020, 257 patients were included in the study. Mean age 57.9 years (SD 10.1), mostly women (98.8%), with children (87.9%) and married (65.4%). According to clinical status 75.5% were non-metastatic and 24.5% metastatic. Protocol compliance was achieved in more than 90% in all questionnaires without differences between metastatic and no metastatic patients. **Conclusions:** This multidimensional protocol enables us to make an integral assessment of the QoL of the patients and their unmet needs, as well as to reflect the patients' concerns, both in early and advanced stages of the disease, complementing currently available assessment methods. In the next future a complied questionnaire with the most challenging questions could be developed to be useful as a routine instrument in clinical practice

Introduction: Breast cancer is considered a chronic disease that has considerable impact on the quality of life (QoL) of the patients. Yet, there is not a totally satisfactory evaluation system reflecting the complexity of the breast cancer condition. Furthermore, most instruments are more addressed towards early-stage disease than advanced disease. **Objective:** To describe the methodology used to measure the QoL of life and coping strategies in a representative group of breast cancer patients including metastatic and non-metastatic patients. **Methodology:** Prospective study including patients with breast cancer at different stages of the disease recruited at the Medical Oncology Department of the Catalan Oncology Institute (ICO) previous informed consent. A protocol of approximately one hour face-to-face interview is run to collect sociodemographic information and

to answer the questions of several QoL scales such as the QLCA-AFex Font, QLQ-C30, QLQ-BR23, HADS, DME, BRCS, MINI-MAC, LOT-R and OE questionnaires, complemented by a semi-structured interview. Results: From June 2017 to March 2020, 257 patients were included in the study. Mean age 57.9 years (SD 10.1), mostly women (98.8%), with children (87.9%) and married (65.4%). According to clinical status 75.5% were non-metastatic and 24.5% metastatic. Protocol compliance was achieved in more than 90% in all questionnaires without differences between metastatic and no metastatic patients. Conclusions: This multidimensional protocol enables us to make an integral assessment of the QoL of the patients and their unmet needs, as well as to reflect the patients' concerns, both in early and advanced stages of the disease, complementing currently available assessment methods. In the next future a completed questionnaire with the most challenging questions could be developed to be useful as a routine instrument in clinical practice

<https://rebiunoda.pro.baratznet.cloud:28443/OpacDiscovery/public/catalog/detail/b2FpOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMjg4NzYyNjk>

Título: Development of a holistic strategy for the assessment of the quality of life in patients with breast cancer in the different stages of the disease [electronic resource]

Editorial: Asociación de Psicooncología de Madrid 2021-03-01

Tipo Audiovisual: Quality of life; breast cancer; advanced disease; protocol; multidimensional approach; unmet needs Calidad de vida; cáncer de mama; enfermedad avanzada; protocolo; enfoque multidimensional; necesidades no atendidas

Variantes del título: Desarrollo de una estrategia holística para la valoración de la calidad de vida en pacientes con cáncer de mama en las distintas etapas de la enfermedad

Documento fuente: Psicooncología; Vol 18 No 1 (2021); 117-136

Nota general: application/pdf

Restricciones de acceso: Open access content. Open access content star

Condiciones de uso y reproducción: Derechos de autor 2021 Psicooncología

Lengua: Spanish

Enlace a fuente de información: Psicooncología; Vol 18 No 1 (2021); 117-136 Psicooncología; Vol. 18 Núm. 1 (2021); 117-136 1988-8287 1696-7240

Otras relaciones: <https://revistas.ucm.es/index.php/PSIC/article/view/74535/4564456556701> /*ref*/Talepasand S, Mahfar F. Relationship between defense mechanisms and the quality of life in women with breast cancer. Int J Cancer Manag 2018;11:1-8. doi: <https://doi.org/10.5812/ijcm.11116> /*ref*/Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Cancer J Clin 2018;68:394-424. doi: <https://doi.org/10.3322/caac.21492> /*ref*/Asociación Española Contra el Cáncer. Pronóstico del Cáncer de Mama: Mortalidad y Esperanza de vida [Internet]. 2018 [citado 20 de noviembre de 2019]. Disponible en: <https://www.aecc.es/es/todo-sobre-cancer/tipos-cancer/cancer-mama/mas-informacion/evolucion-cancer-mama> /*ref*/Izquierdo A, Gispert R, Saladie F, Espinàs JA. Análisis de la incidencia, la supervivencia y la mortalidad según las principales localizaciones tumorales, 1985-2019: cáncer de mama. Med Clínica 2008;131:50-2. doi: [https://doi.org/10.1016/S0025-7753\(08\)76433-9](https://doi.org/10.1016/S0025-7753(08)76433-9) /*ref*/Koch L, Jansen L, Herrmann A, Stegmaier C, Holleczeck B, Singer S, et al. Quality of life in long-term breast cancer survivors - a 10-year longitudinal population-based study. Acta Oncol 2013;52:1119-28. doi: <https://doi.org/10.3109/0284186X.2013.774461> /*ref*/Cardoso F, Costa A, Senkus E, Aapro M, André F, Barrios CH, et al. 3rd ESO-ESMO International Consensus Guidelines for Advanced Breast Cancer (ABC 3). Ann Oncol 2017;28:16-33. doi: <https://doi.org/10.1093/annonc/mdw544> /*ref*/Cardoso F. Metastatic breast cancer patients: The forgotten heroes!. Breast 2009;18:271- 2. doi: <https://doi.org/10.1016/j.breast.2009.09.001> /*ref*/Aziz NM. Cancer survivorship research: State of knowledge, challenges and opportunities. Acta Onco. 2007;46:417-32. doi: <https://doi.org/10.1080/02841860701367878> /*ref*/Costa-Requena G, Rodríguez A, Fernández-Ortega P. Longitudinal assessment of distress and quality of life in the early stages of breast cancer treatment: Breast cancer

and longitudinal distress. *Scand J Caring Sci* 2013;27:77-83. doi: <https://doi.org/10.1111/j.1471-6712.2012.01003.x>

/*ref*/Hernández F, Landero R. Aspectos psicosociales relacionados con la calidad de vida en mujeres con cáncer de mama Psychosocial aspects related to quality of life in women with breast cancer. *Summa Psicológica* 2014;11:99-104 /*ref*/Hagedoorn M, Sanderman R, Bolks HN, Tuinstra J, Coyne JC. Distress in couples coping with cancer: A meta-analysis and critical review of role and gender effects. *Psychol Bull* 2008;134:1-30. doi: <https://doi.org/10.1037/0033-2909.134.1.1> /*ref*/Arraras JI, Illarramendi JJ, Manterola A, Domínguez Á. Progresos en la evaluación de la calidad de vida en el cáncer de mama. El sistema de medida de la EORTC. *Psicooncología* 2007;4:367-84 /*ref*/Falo C, Villar H, Rodríguez E, Mena M, Font A. Calidad de vida en cáncer de mama metastásico de larga evolución: La opinión de los pacientes. *Psicooncología* 2014;11:313-31. doi: <https://doi.org/10.5209/revPSIC.2014.v11.n2-3.47391> /*ref*/Ferlay J, Shin H-R, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010;127:2893-917. doi: <https://doi.org/10.1002/ijc.25516> /*ref*/Pastells S, Font A. Propuesta de un instrumento para la detección del optimismo estratégico en cáncer de mama. *Psicooncología* 2019;16:315-28. doi: <https://doi.org/10.5209/revPSIC.2014.v11.n1.44914> /*ref*/Font A. Valoración de la calidad de vida en pacientes de cáncer. [Tesi Doctoral]. [Bellaterra, Barcelona]: Universitat Autònoma de Barcelona.; 1988 /*ref*/Font A. Cáncer y calidad de vida. *Anu Psicol* 1994;61:41-50 /*ref*/Padierna C, Fernández C, González A. Calidad de vida en pacientes oncológicos terminales, medida con el QL-CA-AFEX. *Psicothema* 2002;14:1-8 /*ref*/Aaronson NK, Ahmedzai S, Bergman B, Bullinger M, Cull A, Duez NJ, et al. The European Organization for Research and Treatment of Cancer QLQ-C30: A quality-of-life instrument for use in international clinical trials in oncology. *JNCI J Natl Cancer Inst* 1993;85:365-76 /*ref*/Sprangers MA, Groenvold M, Arraras JI, Franklin J, te Velde A, Muller M, et al. The European Organization for Research and Treatment of Cancer breast cancer-specific quality-of-life questionnaire module: first results from a three-country field study. *J Clin Oncol* 1996;14:2756-68. doi: <https://doi.org/10.1200/JCO.1996.14.10.2756> /*ref*/Finck C, Barradas S, Zenger M, Hinz A. Quality of life in breast cancer patients: Associations with optimism and social support. *Int J Clin Health Psychol* 2018;18:27-34. doi: <https://doi.org/10.1016/j.ijchp.2017.11.002> /*ref*/Hinz A, Singer S, Brähler E. European reference values for the quality of life questionnaire EORTC QLQ-C30: Results of a German investigation and a summarizing analysis of six European general population normative studies. *Acta Oncol*. 2014;53:958-65. doi: <https://doi.org/10.3109/0284186X.2013.879998> /*ref*/Cerezo O, Oñate-Ocaña LF, Arrieta-Joffe P, González-Lara F, García-Pasquel MJ, Bargalló-Rocha E, et al. Validation of the Mexican Spanish version of the EORTC QLQ C30 and BR23 questionnaires to assess health related quality of life in Mexican women with breast cancer. *Eur J Cancer Care (Engl)*. 2012;21:684-691. doi: <https://doi.org/10.1111/j.1365-2354.2012.01336.x> /*ref*/Arraras JI, Tejedor M, Illaramendi J j., Vera R, Pruja E, Marcos M, et al. The EORTC breast cancer quality questionnaire (QLQ-BR23): A psychometric study with Spanish patients. *Psicol Conduct* 2001;9:81-97 /*ref*/Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand*. 1983;67:361-70. doi: <https://doi.org/10.1111/j.1600-0447.1983.tb09716.x> /*ref*/Terol M, López-Roig S, Rodríguez-Marín J, Gelabert M, Pastor M-Á, Reig MT. Propiedades psicométricas de la Escala Hospitalaria de Ansiedad y Estrés (HAD) en población española. *Ansiedad Estrés* 2007;13:163-76 /*ref*/Maté J, Mateo D, Bayés R, Bernaus M, Casas C, González-Barboteo J, et al. Elaboración y propuesta de un instrumento para la detección de malestar emocional en enfermos al final de la vida. *Psicooncología* 2009;6:507-18 /*ref*/Limonero JT, Mateo D, Maté-Méndez J, González-Barboteo J, Bayés R, Bernaus M, et al. Evaluación de las propiedades psicométricas del cuestionario de Detección de Malestar Emocional (DME) en pacientes oncológicos. *Gac Sanit* 2012;26:145-52. doi: <https://doi.org/10.1016/j.gaceta.2011.07.016> /*ref*/Watson M, Law M, dos Santos M, Greer S, Baruch J, Bliss J. The Mini-MAC: Further development of the Mental Adjustment to Cancer Scale. *J Psychosoc Oncol* 1994;12:33-46. doi: <https://doi.org/10.1300/J077V12N0303> /*ref*/Calderón C, Carmona-Bayonas A, Hernández R, Ghanem I, Castelo B, Martínez de Castro E, et al. Effects of pessimism, depression, fatigue, and pain on functional health-related quality of life in patients with resected non-advanced breast cancer. *Breast* 2019;44:108-12. doi: <https://doi.org/10.1016/j.breast.2019.01.012> /*ref*/Sinclair VG, Wallston KA. The development and psychometric evaluation of the Brief Resilient Coping Scale. *Assessment* 2004;11:94-101. doi: <https://doi.org/10.1177/1073191103258144> /*ref*/Otero JM, Luengo A, Romero E, Gómez JA, Castro C. *Psicología de personalidad. Manual de prácticas*. Barcelona: Ariel Practicum; 1998 /*ref*/Ferrando PJ, Chico E, Tous JM. Propiedades psicométricas del test de optimismo Life Orientation Test. *Psicothema*. 2002;14:673-80 /*ref*/Molenberghs G, Kenward M. Missing data in clinical studies. Chichester, UK: John Wiley and Sons Ltd; 2007 /*ref*/Moons P, Budts W, De Geest S. Critique on the conceptualisation of quality of life: A review and evaluation of different conceptual approaches. *Int J Nurs Stud* 2006;43:891-901. doi: <https://doi.org/10.1016/j.ijnurstu.2006.03.015> /*ref*/Sánchez-Pedraza R, Ballesteros MP,

Anzola JD. Concepto de calidad de vida en mujeres con cáncer. Rev Salud Pública 2010;12:889-902 /*ref*/WHO Quality of Life Assessment Group. Qué calidad de vida? / Grupo de la OMS sobre la calidad de vida. Foro Mund Salud 1996;174):385-7 /*ref*/Rowland JH, Baker F. Introduction: Resilience of cancer survivors across the lifespan. Cancer. 2005;104(S11):2543-8. doi: <https://doi.org/10.1002/cncr.21487> /*ref*/Derogatis LR, Abeloff MD, Melisaratos N. Psychological coping mechanisms and survival time in metastatic breast cancer. JAMA 1979;242: 1504-8. doi: <https://doi.org/10.1001/jama.1979.03300140020016> /*ref*/Scheier MF, Carver CS. Optimism, coping, and health: assessment and implications of generalized outcome expectancies. Health Psychol 1985;4:219-47. doi: <https://doi.org/10.1037/0278-6133.4.3.219> /*ref*/Seligman MEP. Aprenda Optimismo: Haga de la vida una experiencia maravillosa. Barcelona: Clave, 2011 /*ref*/Reed E, Kössler I, Hawthorn J. Quality of life assessments in advanced breast cancer: should there be more consistency?: Quality of life assessment in advanced breast cancer. Eur J Cancer Care (Engl) 2012;21:565-80. doi: <https://doi.org/10.1111/j.1365-2354.2012.01370.x> /*ref*/Irarrázaval ME, Rodríguez PF, Fasce G, Silva FW, Waintrub H, Torres C, et al. Calidad de vida en cáncer de mama: validación del cuestionario BR23 en Chile. Rev Médica Chile 2013;141:723-34 /*ref*/Rosedale M, Fu MR. Confronting the unexpected: temporal, situational, and attributional dimensions of distressing symptom experience for breast cancer survivors. Oncol Nurs Forum 2010;37:E28-33 /*ref*/Caruana EJ, Roman M, Hernández-Sánchez J, Solli P. Longitudinal studies. J Thorac Dis 2015;7:E537-40. doi: <https://doi.org/10.3978/j.issn.2072-1439.2015.10.63>

Baratz Innovación Documental

- Gran Vía, 59 28013 Madrid
- (+34) 91 456 03 60
- informa@baratz.es