



Parents receiving cancer diagnosis: emotional distress and resources to communicate with their children [

Asociación de Psicooncología de Madrid,
2017-10-13

[info:eu-repo/semantics/article](https://doi.org/10.1016/j.annonc.2017.10.013)

[info:eu-repo/semantics/publishedVersion](https://doi.org/10.1016/j.annonc.2017.10.013)

Artículo revisado por pares

Analítica

Objective: To know which resources to communicate the illness to their children and whether these resources are related with emotional distress in cancer patients at diagnosis. **Method:** Sixty patients aged 34-60 years, who have been recently diagnosed and had children less of 18 years old reported their resources, emotional distress and family communication style. Family communication was assessed using the specific subscale of the FAD (McMaster Family Assessment Device). Patients resources were assessed with a specific questionnaire whose items were valued in a 0-10 scale (being 0 nothing and 10 the highest punctuation possible). Emotional distress was assessed with a numeric scale ranging 0-10. **Results:** Resources more available were those related with the management of daily life (mean= 8.3), followed by resources to delegate to others the care of their children. Both resources were available for more than a 80% of patients. However, resources related with telling the children the illness situation, were available to only 55% of patients. Patients with resources to communicate the illness situation to their children reported less emotional distress. **Conclusions:** Parents who were recently diagnosed with cancer have more resources to manage their daily life and to delegate the care of their children to others, than to inform them about the illness and to help them to cope emotionally. Parents with less resources to communicate with their children show a higher level of emotional distress

Objective: To know which resources to communicate the illness to their children and whether these resources are related with emotional distress in cancer patients at diagnosis. **Method:** Sixty patients aged 34-60 years, who have been recently diagnosed and had children less of 18 years old reported their resources, emotional distress and family communication style. Family communication was assessed using the specific subscale of the FAD (McMaster Family Assessment Device). Patients resources were assessed with a specific questionnaire whose items were valued in a 0-10 scale (being 0 nothing and 10 the highest punctuation possible). Emotional distress was assessed with a numeric scale ranging 0-10. **Results:** Resources more available were those related with the management of daily life (mean= 8.3), followed by resources to delegate to others the care of their children. Both resources were available for more than a 80% of patients. However, resources related with telling the children the illness situation, were available to only 55% of patients. Patients with resources to communicate the illness situation to their children reported less emotional distress. **Conclusions:** Parents who were recently diagnosed with cancer have more resources to manage their daily life and to delegate the care of their children to others, than to inform them about the illness and to help them to cope emotionally. Parents with less resources to communicate with their children show a higher level of emotional distress

Título: Parents receiving cancer diagnosis: emotional distress and resources to communicate with their children [electronic resource]

Editorial: Asociación de Psicooncología de Madrid 2017-10-13

Tipo Audiovisual: Cancer communication; psychological resources; ill parents; sons; emotional distress
Comunicación en cáncer; recursos psicológicos; padres enfermos; hijos; malestar emocional

Variantes del título: Padres diagnosticados de cáncer: malestar emocional y recursos para comunicarse con sus hijos meno

Documento fuente: Psicooncología; Vol 14 No 2-3 (2017); 229-240

Nota general: application/pdf

Restricciones de acceso: Open access content. Open access content star

Lengua: Spanish

Enlace a fuente de información: Psicooncología; Vol 14 No 2-3 (2017); 229-240 Psicooncología; Vol. 14 Núm. 2-3 (2017); 229-240 1988-8287 1696-7240

Otras relaciones: <https://revistas.ucm.es/index.php/PSIC/article/view/57082/51518> /*ref*/Gamo Medina E, Pazos Pezzi P. El duelo y las etapas de la vida. Rev Asoc Esp Neuropsiq 2009;29: 455-69 /*ref*/Kroen WC. Cómo ayudar a los niños a afrontar la pérdida de un ser querido - Ed. Oniro S.A. Barcelona. 2002 /*ref*/Lacasta M.A. Soler M.C. El duelo: prevención y tratamiento del duelo patológico. Cuidados después de la muerte Manual SEOM de Cuidados Continuos. 713-30. Dispublic S.L. 2004; Madrid /*ref*/Fearnley R. Death of a parent and the children's experience: Don't ignore the elephant in the room. J Interprof Care. 2010; 24:450-9. doi: 10.3109/13561820903274871 /*ref*/Rodríguez, A. M. Els nens i adolescents davant el càncer avançat o terminal dels pares. (Tesi Doctoral). Barcelona: Universitat Autònoma de Barcelona. 2013 /*ref*/Inhestern y Bultmann, J.C. Understanding parenting concerns in cancer survivors with minor and young-adult children. J Psychosom Res 2016; 87: 1-6. doi: 10.1016/j.jpsychores.2016.05.008 /*ref*/Semple C, McCaughan E. Family life when a parent is diagnosed with cancer: impact of a psychosocial intervention for young children. Eur J Cancer Care 2012; 22: 219-31. doi: 10.1111/ecc.12018 /*ref*/Lewis F, Casey S, Brandt P, Shands ME, Zahlis EH. The enhancing connections program: pilot study of a cognitive-behavioral intervention for mothers and children affected by breast càncer. Psychooncology 2005; 15: 486-97 /*ref*/Fuentes S, Blasco T. Preocupaciones de los pacientes con càncer de mama que reciben atención psicooncológica. Psicooncología 2010;7: 51-60 /*ref*/Fuentes S, Blasco T. Factores predictores de la adaptación a la enfermedad en pacientes recién diagnosticadas de càncer de mama que acuden a un servicio de psicooncología. Anal Psicol 2012; 28:736-42. doi:10.6018/analesps.28.3.156031 /*ref*/Arber,A. 'Think Family Project' International Conference on Cancer Nursing Vancouver, BC Canada; 2015 /*ref*/Sordo T, Rubio I, Peralta M, Ángeles M, Ospina L. Cómo se lo digo a mi hijo/a? Psicooncología 2014; 11(supl. 1):1-83 /*ref*/Thastum M, Watson M, Kiembarcher C, Piha J, Steck B, Zachariae, et al. Prevalence and predictors of emotional and behavioral functioning of children where a parent has a cancer: a multinational study. Cancer. 2009; 115: 4030-9. doi: 10.1002/cncr.24449 /*ref*/Huizinga G, Visser A, Zelders-Steyn Y, Teule J, Reijneveld S, Roodbol P. Psychological impact of having a parent with cancer. Eur J Cancer 2011;47: 239-46. doi: 10.1016/S0959-8049(11)70170-8 /*ref*/Krattenmacher T, Kuhne F, Ernst J, Bergelt C, Romer G, Moller, B. Parental cancer: factors associated with children's psychosocial adjustment a systematic review. J Psychosom Res 2012; 72: 344-56. doi: 10.1016/j.jpsychores.2012.01.011 /*ref*/Krattenmacher T, Kühne F, Führer D, Beierlein V, Brähler E, Resch F, et al. Coping skills and mental health status in adolescents when a parent has cancer: A multicenter and multi-perspective study. J Psychosom Res 2013; 74, 252-9. doi: 10.1016/j.jpsychores.2012.10.003 /*ref*/Phillip, F. Adolescents living with a parent with advanced cancer: a review of the literature. Psychooncology 2014;23:1323-39. doi: 10.1002/pon.3570 /*ref*/Ellis SJ, Wakefield CE, Burns M. Supporting children facing a parent's cancer diagnosis: a systematic review of children's psychosocial needs and existint interventions. Eur J Cancer Care 2017; 26 (1). doi: 10.1111/ecc.12432 /*ref*/Mesquida V, Seijas R, Rodríguez M. Los niños ante la pérdida de uno de los

progenitores: revisión de pautas de comunicación eficaces. *Psicooncología* 2015;12:417-29. doi: 10.5209/revPSIC.2015.v12.n2-3.51019 /*ref*/*Britz R. El álbum de mamá. Barcelona . Novartis Oncology, 2014 /*ref*/*Elizegi, Rebeka. Una flor de recanvi per a la mare. Barcelona: Editorial Takatuka, 2011 /*ref*/*Fuentes S, Giralt M. Qué te ocurre..., mamá? Barcelona: Ed. Acción Médica, 2006 /*ref*/*Epstein N, Baldwin L, Bishop D. The McMaster Family Assessment Device. *J Marital Fam Ther* 1983;9:171-80. doi: 10.1111/j.1752-0606.1983.tb01497.x /*ref*/* /Fuentes, S. Comunicació en càncer: Anàlisi de les preocupacions i dels recursos dels pares recién diagnosticats al comunicar la malaltia als seus fills. (Tesi Doctoral). Barcelona: Universitat Autònoma de Barcelona. 2015 /*ref*/* /Gessler S, Low J, Daniells E, Williams R, Brough V, Tookman A, et al. Screening for distress in cancer patients: is the distress thermometer a valid measure in the UK and does it measure change over time?. A prospective validation study. *Psychooncology* 2008; 17:538-47. doi: 10.1002/pon.1273 /*ref*/*Priede A, Torres M, Hoyuela F, González-Blanch C. El termómetro del distress como prueba de cribado de ansiedad y depresión en pacientes oncológicos recién diagnosticados. *Psicooncología* 2014;11:31-43. doi: 10.5209/revPSIC.2014.v11.n1.44915 /*ref*/* /Zwahlen D, Hagenbuck N, Carley M, Recklitis C, Buchi S. Screening cancer patients' families with the distress thermometer (DT): a validation study. *Psychooncology* 2008;17:959-66. doi: 10.1002/pon.1320 /*ref*/*Kim S, Ko Y, Jun E. The impact of breast cancer on mother-child relationship in Korea. *Psychooncology* 2012;21 640-6. doi: 10.1002/pon.1941 /*ref*/*Bugge K, Helseth S, Darbyshire P. Parent's experiences of a Family Support Program when a parent has incurable cancer. *J Clin Nurs* 2009; 8:3480-8. doi: 10.1111/j.1365-2702.2009.02871.x. /*ref*/* /Romer G, Saha R, Haagen M, Pott M, Baldus C, Bergelt C. Lessons learned in the implementation of an innovative consultation and liaison service for children of cancer patients in various hospital settings. *Psychooncology* 2007;16:138-48. doi: 10.1002/pon.1105 /*ref*/*Fasciano K, Berman H, Moore C, DeFrino B, Jameson R, Kennedy S, et al. When a parent has a cancer: a community based program for school personnel. *Psychooncology* 2007;16:158-67. doi:10.1002/pon.1148 /*ref*/*Huang X, O'Connor M, Lee S. School-aged and adolescent children's experience when a parent has non-terminal cancer: a systematic review and meta-synthesis of qualitative studies. *Psychooncology* 2014;23:493-506. doi: 10.1002/pon.3457 /*ref*/*Grenklo T, Kreicbergs U, Valdimarsdóttir U, Nyberg T, Steineck G, Fürst C. Self-injury in youths who lost a parent to cancer: nationwide study of the impact of family-related and health-care-related factors. *Psychooncology* 2014;23:989-97. doi: 10.1002/pon.3515 /*ref*/*Kopchak D, Burke C. Interaction patterns between parents with advanced cancer and their adolescent children. *Psychooncology* 2011; 20:1108-15. doi.10.1002/pon.1831 /*ref*/*Muñoz MJ, Bernal P. Abordaje de la ansiedad en pacientes diagnosticados de càncer. *Psicooncología* 2016;13:227-48. doi:10.5209/PSIC.54434 /*ref*/*Rodríguez AM. Cómo explicar a los niños y adolescentes el càncer avanzado o terminal de sus padres? *Psicooncología* 2015; 12:171-82. doi: 10.5209/revPSIC.2015.v12.n1.48911 /*ref*/*Vila L, Font A, Caba L. Estado emocional en mujeres con càncer de mama: variación experimentada después de una sesión psico-oncológica basada en el counselling y la psicología positiva. *Psicooncología*. 2016; 13: 205-25. doi:10.5209/PSIC.54433

Baratz Innovación Documental

- Gran Vía, 59 28013 Madrid
- (+34) 91 456 03 60
- informa@baratz.es