



Semi-structured interview for the evaluation of oncological disease awareness E-CEO. Reliability study on a sample of patients with advanced disease [

Asociación de Psicooncología de Madrid,
2019-04-08

[info:eu-repo/semantics/article](https://doi.org/10.1016/j.baratz.2019.04.008)

[info:eu-repo/semantics/publishedVersion](https://doi.org/10.1016/j.baratz.2019.04.008)

Artículo revisado por pares

Analítica

Objective: Communication between cancer patients and health professionals is considered one of the most important aspects of care, especially in advanced stages of the disease. Disease awareness, consists of a dynamic and evolving process over time, influenced by the quantity and quality of information and significance that each individual assigns. In this paper, the process for the design, validation and reliable calculation of a semi-structured interview (Interview on the Consciousness of Oncological Disease E-CEO) is described. This process is sparingly intrusive, non-disclosing of information and makes it possible to place an individual along a continuum according to their degree of perception of disease severity. **Method:** In an initial phase, an open interview was conducted with 15 patients and a list was prepared of key expressions, which were then grouped into five categories. In a second phase, the semi-structured interview model was applied to 42 patients admitted with a diagnosis of advanced oncological disease (clinical stage IV) in the Oncology Hospitalization Unit of the Hospital of A Coruña. **Results:** The reliability obtained among observers shows that it is a high-quality instrument ($Kappa = .70$). The data reflects a wide discrepancy between the health status of the patients and their own perception. More than 50% consider their illness "controlled". **Conclusions:** This is a simple application that will allow future in-dept research into the study of information integration processes for these types of patients

Objective: Communication between cancer patients and health professionals is considered one of the most important aspects of care, especially in advanced stages of the disease. Disease awareness, consists of a dynamic and evolving process over time, influenced by the quantity and quality of information and significance that each individual assigns. In this paper, the process for the design, validation and reliable calculation of a semi-structured interview (Interview on the Consciousness of Oncological Disease E-CEO) is described. This process is sparingly intrusive, non-disclosing of information and makes it possible to place an individual along a continuum according to their degree of perception of disease severity. **Method:** In an initial phase, an open interview was conducted with 15 patients and a list was prepared of key expressions, which were then grouped into five categories. In a second phase, the semi-structured interview model was applied to 42 patients admitted

with a diagnosis of advanced oncological disease (clinical stage IV) in the Oncology Hospitalization Unit of the Hospital of A Coruña. Results: The reliability obtained among observers shows that it is a high-quality instrument (Kappa = .70). The data reflects a wide discrepancy between the health status of the patients and their own perception. More than 50% consider their illness "controlled". Conclusions: This is a simple application that will allow future in-dept research into the study of information integration processes for these types of patients

<https://rebiunoda.pro.baratznet.cloud:28443/OpacDiscovery/public/catalog/detail/b2FpOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMjkwOTY4MTU>

Título: Semi-structured interview for the evaluation of oncological disease awareness E-CEO. Reliability study on a sample of patients with advanced disease [electronic resource]

Editorial: Asociación de Psicooncología de Madrid 2019-04-08

Tipo Audiovisual: cancer; disease awareness; semi-structured interview cáncer; conciencia de enfermedad; entrevista semiestructurada

Variantes del título: Entrevista semi-estructurada para la evaluación de la conciencia de enfermedad oncológica E-CEO. Estudio de fiabilidad sobre una muestra de pacientes con enfermedad avanzada

Documento fuente: Psicooncología; Vol 16 No 1 (2019); 11-23

Nota general: application/pdf

Restricciones de acceso: Open access content. Open access content star

Lengua: Spanish

Enlace a fuente de información: Psicooncología; Vol 16 No 1 (2019); 11-23 Psicooncología; Vol. 16 Núm. 1 (2019); 11-23 1988-8287 1696-7240

Otras relaciones: <https://revistas.ucm.es/index.php/PSIC/article/view/63645/4564456549463> /*ref*/Andruccioli J, Montesi A, Di Leo S, Sarti D, Turci P, Pittureri C, et al . Illness awareness in hospice: application of a semi-structured interview. Am J Hosp Palliat Care 2009;26:384-91 /*ref*/Andruccioli J, Montesi A, Raffaeli W, Monterubbianesi MC, Turci P, Pittureri C, et al. Illness awareness of patients in hospice: psychological evaluation and perception of family members and medical staff. J Palliat Med 2007;10:741-8 /*ref*/Applebaum AJ, Kolva EA, Kulikowski JR, Jacobs JD, DeRosa A, Lichtenthal WG, Olden ME, Rosenfeld B, Breitbart W. Conceptualizing prognostic awareness in advanced cancer: a systematic review. J Health Psychol 2014;19:1103-19 /*ref*/Burns CM, Broom DH, Smith WT, Dear K, Craft PS. Fluctuating awareness of treatment goals among patients and their caregivers: A longitudinal study of a dynamic process. Support Care in Cancer. 2007;15:187-96 /*ref*/Camprubi N, Almela A, Garre-Olmo J. Propiedades psicométricas de la validación española de la Insight Scale. Actas Españolas de Psiquiatría 2008; 36: 323-30 /*ref*/Caruso A, Di Francesco B, Pugliese P, Cinanni V, Corlito A. Information and awareness of diagnosis and progression of cancer in adult and elderly cancer patients. Tumori 2000;86:199-203 /*ref*/Centeno-Cortes C, Nuñez-Olarte JM. Estudios sobre la comunicación del diagnóstico del cáncer en España. Med Clin 1998; 110:744-50 /*ref*/Centeno-Cortes C, Nuñez-Olarte JM. Questioning diagnosis disclosure in terminal cancer patients: A prospective study evaluating patients responses. Palliat Med 1994;8:39-44 /*ref*/Chominov HM, Tataryn DJ, Wilson KG, Enns M, Lander S. Prognosis awareness and the terminally ill. Psychosomatic. 2000; 41: 500-4 /*ref*/Corli O, Apolone G, Pizzuto M, Cesaris L, Cozzolino A, Orsi L, et al. Illness awareness in terminal cancer patients: an Italian study. Palliat Med 2009;23:354-9 /*ref*/Costantini A, Grassi L, Picardi A, Brunetti S, Caruso R, Nanni MG, et al. Awareness of cancer, satisfaction with care, emotional distress, and adjustment to illness: an Italian multicenter study. Psychooncology 2015;24:1088-96. doi: 10.1002/pon.3768 /*ref*/Craft PS, Burns CM, Smith WT, Broom DH. Knowledge of treatment intent among patients with advanced cancer: a longitudinal study. Eur J Cancer Care 2005;14:417-25 /*ref*/Degner LF, Kristjanson LJ, Bowman D, Sloan JA, Carriere KC, O'Neil J, et al. Information needs and decisional preferences in women with breast cancer. JAMA. 1997; 277:1485-92 /*ref*/Epstein RM, Street RL. [Internet] Patient-Centered Communication in Cancer Care: Promoting Healing and Reducing Suffering. Bethesda (MD): National Cancer Institute. 2007 /*ref*/Estapé J, Palombo H, Hernandez E, Daniels M, Estapé T, Grau JJ, et al. Cancer diagnosis

disclosure in a Spanish hospital. *Ann Oncol* 1992; 3: 451-4 /ref*/Fallowfield L, Jenkins V. Communicating sad, bad, and difficult news in medicine. *Lancet* 2004;363:312-9 /ref*/Field D, Copp G. Communication and awareness about dying in the 1990s. *Palliat Med* 1999; 13: 459-68 /ref*/Friis LS, Elverdam B, Schmidt KG. The patient's perspective. A qualitative study of acute myeloid leukaemia patients' need for information and their information-seeking behaviour. *Support Care Cancer*. 2003;11:162-70 /ref*/Gattellari M, Butow PN, Tattersall MH, et al. Misunderstanding in cancer patients: Why shoot the messenger? *Ann Oncol* 1999;10:39-46. Gordon EJ, Daugherty CK. 'Hitting you over the head': oncologists' disclosure of prognosis to advanced cancer patients. *Bioethics* 2003;17:142-68 /ref*/Hagerty RG, Butow PN, Ellis PM, Dimitry S, Tattersall MH. Communicating prognosis in cancer care: a systematic review of the literature. *Ann Oncol* 2005;16:1005-53 /ref*/Helft PR, Hlubocky F, Wen M, Daugherty CK. Associations among awareness of prognosis, hopefulness, and coping in patients with advanced cancer participating in phase I clinical trials. *Support Care Cancer*. 2003;11:644-51 /ref*/Iconomou G, Viha A, Koutra A, Vagenakis AG, Kalofonos HP. Information needs and awareness of diagnosis in patients with cancer receiving chemotherapy: A report from Greece. *Palliat Med* 2002;16:315-21 /ref*/Innes S, Payne S. Advanced cancer patients' prognostic information preferences: a review. *Palliat Med* 2009; 23: 29-39 /ref*/Johnston G, Abraham C. Managing awareness: Negotiating and coping with a terminal prognosis. *Inter J Palliat Nurs* 2000; 6: 485-94 /ref*/Kaplowitz SA, Campo S, Chiu WT. Cancer patients' desires for communication of prognosis information. *Health Commun* 2002; 14:221-41 /ref*/Karnofsky DA, Burchenal JH. The clinical evaluation of chemotherapeutic agents in cancer. En: CM Macleod (ed.). *Evaluation of chemotherapeutic agents*. New York: Columbia University Press, 1949; p. 191-205 /ref*/Landis J, Koch G. The measurement of observer agreement for categorical data. *Biometrics* 1977; 33:159-74 /ref*/Leydon GM, Boulton M, Moynihan C, Jones A, Mossman J, Boudioni M, et al. Cancer patients' information needs and information seeking behaviour: in depth interview study. *BMJ* 2000; 320:909-13 /ref*/Lynn MR. Determination and quantification of content validity. *Nurs Res* 1986;35:382-5 /ref*/Malouf T, Langdon R, Taylor A. The Insight Interview: a new tool for measuring deficits in awareness after traumatic brain injury. *Brain Injury* 2014; 28: 1523-41 /ref*/Martos-Gutiérrez JA, Montoro-Lorite M, Mundet-Pons L. Quality of diagnosis information given to terminal cancer patients. *Enferm Clin* 2016; 26:344-50. doi: 10.1016/j.enfcli.2016.07.006 /ref*/Morasso G, Alberisio A, Capelli M, Rossi C, Baracco G, Costantini M. Illness awareness in cancer patients: a conceptual framework and a preliminary classification hypothesis. *Psychooncology* 1997; 6: 212-7 /ref*/Muthu Kumar D, Symonds RP, Sundar S, Ibrahim K, Savelyich BS, Miller E. Information needs of Asian and white British cancer patients and their families in Leicestershire: A cross-sectional survey. *Br J Cancer* 2004; 90:1474-8 /ref*/Nord C, Mykletun A, Fossa SD. Cancer patients awareness about their diagnosis: A population-based study. *J Public Health Med* 2003;25: 313-7 /ref*/Nuñez Olarte JM. Información al paciente en situación terminal: diferencias entre las publicaciones anglosajonas y la experiencia diaria en España. *Med Paliat* 2013; 21: 113- 20 /ref*/Pimentel FL, Ferreira JS, Vila Real M, Mesquita NF, Maia-Goncalves JP. Quantity and quality of information desired by Portuguese cancer patients. *Support Care Cancer* 1999; 7:407-12 /ref*/Pronzato P, Bertelli G, Losardo P, Landucci M. What do advanced cancer patients know of their disease? A report from Italy. *Support Care Cancer* 1994; 2: 242-4 /ref*/Seo M, Tamura K, Shijo H, Morioka E, Ikegame C, Hirasako K. Telling the diagnosis to cancer patients in Japan: attitude and perception of patients, physicians and nurses. *Palliat Med* 2000; 14:105-10. Steinhauser KE, Christakis NA, Clipp EC, McNeilly M, Grambow S, Parker J, et al. Preparing for the end of life: preferences of patients, families, physicians, and other care providers. *J Pain Symptom Manage* 2001;22:727-37 /ref*/Stewart M, Brown JB, Boon H, Galajda J, Meredith L, Sangster M. Evidence on patient-doctor communication. *Cancer Prev Control* 1999; 3:25-30 /ref*/Watson M, Greer S, Blake S, Shrapnell K. Reaction to a diagnosis of breast cancer: relationship between denial, delay, and rates of psychological morbidity. *Cancer* 1984; 53: 2008-12 /ref*/Zachariae R, Pedersen CG, Jensen AB, Ehrnrooth E, Rossen PB, Maase H. Association of perceived physician communication style with patient satisfaction, distress, cancer-related self-efficacy, and perceived control over the disease. *Br J Cancer* 2003; 88: 658-65

Baratz Innovación Documental

- Gran Vía, 59 28013 Madrid
- (+34) 91 456 03 60
- informa@baratz.es

