



Aislamiento de Plesiomonas shigelloides y Aeromonas veronii biotipo sobria en heces de lobo marino común sudamericano, Otaria flavescens (Shaw, 1800) [

Universidad de Valparaiso: Facultad de Ciencias del Mar, 2009

text (article)

Analítica

Plesiomonas shigelloides and Aeromonas spp. are Gram negative bacteria vastly distributed in the environment, being isolated from aquatic ecosystems and terrestrial and marine animals. The South American sea lion (*Otaria flavescens*) is the most frequent marine mammal of the Chilean coasts, living in beaches, rocks or coastline of rivers. In this work we determined the isolation frequency of *P. shigelloides* and *A. veronii* biotype *sobria* in fecal samples of South American sea lions belonging to a colony established at the urban South coastline of Valdivia River, southern Chile. From the 30 samples under study, *P. shigelloides* was isolated in 27 (90.0%) and *A. veronii* biotype *sobria* in 17 (56.6%). To our knowledge, this is the first report of the isolation of *P. shigelloides* and *Aeromonas* spp. from South American sea lions in Chile. However, further studies are needed to clarify if these bacteria play any role in producing disease, or are merely commensals, in these marine mammals

<https://rebiunoda.pro.baratznet.cloud:28443/OpacDiscovery/public/catalog/detail/b2FpOmNlbgVicmF0aW9uOmVzLmJhemF0ei5yZW4vMzA3MTMzMjM>

Título: Aislamiento de Plesiomonas shigelloides y Aeromonas veronii biotipo sobria en heces de lobo marino común sudamericano, Otaria flavescens (Shaw, 1800) [electronic resource]

Editorial: Universidad de Valparaiso: Facultad de Ciencias del Mar 2009

Tipo Audiovisual: Aeromonadaceae Enterobacteriaceae intestinal microbiota marine mammals reservoir

Documento fuente: Revista de biología marina y oceanografía, ISSN 0717-3326, Vol. 44, N°. 3, 2009, pags. 763-765

Nota general: application/pdf

Restricciones de acceso: Open access content. Open access content star

Condiciones de uso y reproducción: LICENCIA DE USO: Los documentos a texto completo incluidos en Dialnet son de acceso libre y propiedad de sus autores y/o editores. Por tanto, cualquier acto de reproducción, distribución, comunicación pública y/o transformación total o parcial requiere el consentimiento expreso y escrito de aquéllos. Cualquier enlace al texto completo de estos documentos deberá hacerse a través de la URL oficial de éstos en Dialnet. Más información: <https://dialnet.unirioja.es/info/derechosOAI> | INTELLECTUAL PROPERTY RIGHTS STATEMENT: Full text documents hosted by Dialnet are protected by copyright and/or related rights. This digital object is accessible without charge, but its use is subject to the licensing conditions set by its authors or editors. Unless expressly stated otherwise in the licensing conditions, you are free to linking, browsing, printing and making a copy for your own personal purposes. All other acts of reproduction and communication to the public are subject to the licensing conditions expressed by editors and authors and require consent from them. Any link to this document should be made using its official URL in Dialnet. More info: <https://dialnet.unirioja.es/info/derechosOAI>

Lengua: Spanish

Enlace a fuente de información: Revista de biología marina y oceanografía, ISSN 0717-3326, Vol. 44, Nº. 3, 2009, pags. 763-765

Baratz Innovación Documental

- Gran Vía, 59 28013 Madrid
- (+34) 91 456 03 60
- informa@baratz.es