

Biofloc technology on the zootechnical performance of tilapia: effect of strain and stocking density [

2017

text (article)

Analítica

The necessity of new technologies for the culture of aquatic organisms is becoming more evident, and thus, the biofloc system, a technic created to perform zero or minimum water exchange, starts to become more useful not only in shrimp culture, but also in finfish culture. This research purpose was to evaluate the growth and survival of Nile tilapia or "gray" (Oreochromis niloticus - GIFT strain) and red tilapia "Red Florida" strain (Oreochromis sp.) in biofloc system stocked in different densities (400 e 800/m3) in brackish water (8‰). Sixteen 26L plastic bins (19L usefull) were used and 192 fingerlings (3,06 " 0,2 g) were stocked. A factorial experimental design was adopted (strain vs density) and a "macrocosm-microcosm" device system. For42 days, the animals were fed on a 45% CP commercial diet, three times per day. Water quality parameters and microbial community were monitored. At the end, proximate analysis of biofloc biomass was performed. The results suggested that Nile tilapia was the most appropriate for this phase (3-20 g) in biofloc system in brackish water. Furthermore, no negative effect was observed when 800 fish/m2 stocking density was considered

The necessity of new technologies for the culture of aquatic organisms is becoming more evident, and thus, the biofloc system, a technic created to perform zero or minimum water exchange, starts to become more useful not only in shrimp culture, but also in finfish culture. This research purpose was to evaluate the growth and survival of Nile tilapia or "gray" (Oreochromis niloticus - GIFT strain) and red tilapia "Red Florida" strain (Oreochromis sp.) in biofloc system stocked in different densities (400 e 800/m3) in brackish water (8‰). Sixteen 26L plastic bins (19L usefull) were used and 192 fingerlings (3,06 " 0,2 g) were stocked. A factorial experimental design was adopted (strain vs density) and a "macrocosm-microcosm" device system. For42 days, the animals were fed on a 45% CP commercial diet, three times per day. Water quality parameters and microbial community were monitored. At the end, proximate analysis of biofloc biomass was performed. The results suggested that Nile tilapia was the most appropriate for this phase (3-20 g) in biofloc system in brackish water. Furthermore, no negative effect was observed when 800 fish/m2 stocking density was considered

https://rebiunoda.pro.baratznet.cloud: 28443/OpacDiscovery/public/catalog/detail/b2FpOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzEyNDMwOTkwicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzEyNDMwOTkwicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzEyNDMwOTkwicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzEyNDMwOTkwicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzEyNDMwOTkwicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzEyNDMwOTkwicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzEyNDMwOTkwicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzEyNDMwOTkwicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzEyNDMwOTkwicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzEyNDMwOTkwicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzEyNDMwOTkwicmF0ei5yZW4vMzEyNDMwoTkwicmF0ei5yZW4vMzEyNDMwoTkwicmF0ei5yZW4vMzEyNDMwoTkwicmF0ei5yZW4vMzEyNDMwoTkwicmF0ei5yZW4vMzEyNDMwoTkwicmF0ei5yZW4vMzEyNDMwoTkwicmF0ei5yZW4vMzEyNDMwoTkwicmF0ei5yZW4vMzEyNDWataWF0ei5yZW4vMzEyN0faeF0ei5yZW4vMzEyNDWataWF0ei5yZW4vMzEyNDWataWF0ei5yZW4vMzEyNDWataWF0ei5yZW4vMzEyN0faeF0ei5yZW4vMzEyNDWataWF0ei5yZW4vMzEyNDWataWF0ei5yZW4vMzEyNDWataWF0ei5yZW4vMzEyNDWataWF0ei5yZW4vMzEyNDWataWF0ei5yZW4vMzEyNDWataWF0ei5yZW4vMzEyNDWataWF0ei5yZW4vMzEyNDWataWF0ei5yZW4vMzEyNDWataWF0ei5yZW4wAtaWF0ei5yZW4wAtaWF0ei5yZW4wAtaWF0ei5yZW4wAtaWF0ei5yZW4wAtaWF0ei5yZW4wAtaWF0ei5yZW4wAtaWF0ei5yF0ei5yZW4wAtaWF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5yF0ei5

Título: Biofloc technology on the zootechnical performance of tilapia: effect of strain and stocking density electronic resource]

Editorial: 2017

Tipo Audiovisual: Oreochromis Lineage Performance BFT Planktonic community Nursery Oreochromis Linhagem Crescimento Bft Comunidade planctônica Berçário

Documento fuente: Archivos de zootecnia, ISSN 0004-0592, Vol. 66, Nº. 254, 2017, pags. 229-235

Nota general: application/pdf

Restricciones de acceso: Open access content. Open access content star

Condiciones de uso y reproducción: LICENCIA DE USO: Los documentos a texto completo incluidos en Dialnet son de acceso libre y propiedad de sus autores y/o editores. Por tanto, cualquier acto de reproducción, distribución, comunicación pública y/o transformación total o parcial requiere el consentimiento expreso y escrito de aquéllos. Cualquier enlace al texto completo de estos documentos deberá hacerse a través de la URL oficial de éstos en Dialnet. Más información: https://dialnet.unirioja.es/info/derechosOAI | INTELLECTUAL PROPERTY RIGHTS STATEMENT: Full text documents hosted by Dialnet are protected by copyright and/or related rights. This digital object is accessible without charge, but its use is subject to the licensing conditions set by its authors or editors. Unless expressly stated otherwise in the licensing conditions, you are free to linking, browsing, printing and making a copy for your own personal purposes. All other acts of reproduction and communication to the public are subject to the licensing conditions expressed by editors and authors and require consent from them. Any link to this document should be made using its official URL in Dialnet. More info: https://dialnet.unirioja.es/info/derechosOAI

Lengua: English

Enlace a fuente de información: Archivos de zootecnia, ISSN 0004-0592, Vol. 66, Nº. 254, 2017, pags. 229-235

Baratz Innovación Documental

- Gran Vía, 59 28013 Madrid
- (+34) 91 456 03 60
- informa@baratz.es