

Aprovechamiento de metabolitos nitrogenados del cultivo de tilapia en un sistema acuapónico [

2016

text (article)

Analítica

During tilapia production cycle (six months), mushrooms, green fodder and cucurbits were conducted in an aquaponic system to evaluate their efficiency in the removal and use of nitrogenous waste. For the study, a recirculation system of three circular tanks of 7mp3(B was used, where 64 tilapias per mp3(B (0.35 " 0.18 g) were stocked. During the first phase, three mushroom cultivation cycles of 20 d were developed in two systems: Aquaponic and traditional irrigation. Subsequently, six 12 d of growth cycles for green fodders (yellow and white maize, wheat and sorghum), and finally one 45 d cucumber and watermelon bicultivation cycles were done. The total cumulative mushroom production in the aquaponic treatment during the three cycles was 16 % higher than the obtained in the traditional treatment. Most of forage production in total fresh biomass was obtained with wheat (47.55 kg), however The higher length in the stems (27 " 2.20 cm) and the greater nitrogen metabolite removal efficiency were achieved with white maize. The cucumber and watermelon plants produced fruits until the second month of culture with total yield of 27.15 kg and 4.84 kg, respectively, and also they are the most efficient to leverage metabolites versus mushrooms and fodder. With the above results, it can be concluded, that during the tilapia production, it is possible to grow various vegetables, and take advantage of nitrogenous metabolites by aquaponic practices

During tilapia production cycle (six months), mushrooms, green fodder and cucurbits were conducted in an aquaponic system to evaluate their efficiency in the removal and use of nitrogenous waste. For the study, a recirculation system of three circular tanks of 7mp3(B was used, where 64 tilapias per mp3(B (0.35 " 0.18 g) were stocked. During the first phase, three mushroom cultivation cycles of 20 d were developed in two systems: Aquaponic and traditional irrigation. Subsequently, six 12 d of growth cycles for green fodders (yellow and white maize, wheat and sorghum), and finally one 45 d cucumber and watermelon bicultivation cycles were done. The total cumulative mushroom production in the aquaponic treatment during the three cycles was 16 % higher than the obtained in the traditional treatment. Most of forage production in total fresh biomass was obtained with wheat (47.55 kg), however The higher length in the stems (27 " 2.20 cm) and the greater nitrogen metabolite removal efficiency were achieved with white maize. The cucumber and watermelon plants produced fruits until the second month of culture with total yield of 27.15 kg and 4.84 kg, respectively, and also they are the most efficient to leverage metabolites versus mushrooms and fodder. With the above results, it can be concluded, that during the tilapia production, it is possible to grow various vegetables, and take advantage of nitrogenous metabolites by aquaponic practices

Título: Aprovechamiento de metabolitos nitrogenados del cultivo de tilapia en un sistema acuapónico electronic resource]

Editorial: 2016

Tipo Audiovisual: Acuacultura acuaponía forraje de maíz sistemas integrados sustentabilidad Aquaculture aquaponic maize fodder integrated system sustainability

Documento fuente: Ecosistemas y Recursos Agropecuarios, ISSN 2007-901X, null 3, No. 7, 2016, pags. 63-73

Nota general: application/pdf

Restricciones de acceso: Open access content. Open access content star

Condiciones de uso y reproducción: LICENCIA DE USO: Los documentos a texto completo incluidos en Dialnet son de acceso libre y propiedad de sus autores y/o editores. Por tanto, cualquier acto de reproducción, distribución, comunicación pública y/o transformación total o parcial requiere el consentimiento expreso y escrito de aquéllos. Cualquier enlace al texto completo de estos documentos deberá hacerse a través de la URL oficial de éstos en Dialnet. Más información: https://dialnet.unirioja.es/info/derechosOAI | INTELLECTUAL PROPERTY RIGHTS STATEMENT: Full text documents hosted by Dialnet are protected by copyright and/or related rights. This digital object is accessible without charge, but its use is subject to the licensing conditions set by its authors or editors. Unless expressly stated otherwise in the licensing conditions, you are free to linking, browsing, printing and making a copy for your own personal purposes. All other acts of reproduction and communication to the public are subject to the licensing conditions expressed by editors and authors and require consent from them. Any link to this document should be made using its official URL in Dialnet. More info: https://dialnet.unirioja.es/info/derechosOAI

Lengua: Spanish

Enlace a fuente de información: Ecosistemas y Recursos Agropecuarios, ISSN 2007-901X, null 3, N°. 7, 2016, pags. 63-73

Baratz Innovación Documental

- Gran Vía, 59 28013 Madrid
- (+34) 91 456 03 60
- informa@baratz.es