



# Prevalencia y hallazgos radiográficos en conejos, cuyos y chinchillas diagnosticados con maloclusión en el Hospital Veterinario de Especialidades en Fauna Silvestre y Etología Clínica de la UNAM, México [

2020

text (article)

Analítica

Lagomorphs and hystricomorphs, as guinea pigs and chinchillas, have common dental characteristics, such as the elodont and hypsodont teeth. These peculiarities predispose to the presentation of malocclusion due to the consumption of inadequate, non-abrasive diets. The purpose of this work is to identify the prevalence of malocclusion in each case. In addition, it aims to make an objective evaluation of the radiological studies of the specimens, based on anatomical reference lines, to determine the main radiographic findings according to the degree of malocclusion clinically diagnosed. 30% of the individuals evaluated presented malocclusion problems being 42.86% rabbits, 16.6% guinea pigs and 23% chinchillas. There is no evidence of significant differences in the presentation by sex. In rabbits, the diagnosis of the specimens from the group I and II were more common, while in guinea pigs they were II and III. Grade I and II were the most diagnosed in rabbits and guinea pigs. In the radiographs, the most common lesions are the overgrowth of the maxillary and mandibular roots, elongation of the incisors, loss of the occlusal surface and deformation of the mandibular roots. A method of co-rection for LL radiographic projections with evident rotation is proposed to maintain the objectivity of the radiographic evaluation

Lagomorphs and hystricomorphs, as guinea pigs and chinchillas, have common dental characteristics, such as the elodont and hypsodont teeth. These peculiarities predispose to the presentation of malocclusion due to the consumption of inadequate, non-abrasive diets. The purpose of this work is to identify the prevalence of malocclusion in each case. In addition, it aims to make an objective evaluation of the radiological studies of the specimens, based on anatomical reference lines, to determine the main radiographic findings according to the degree of malocclusion clinically diagnosed. 30% of the individuals evaluated presented malocclusion problems being 42.86% rabbits, 16.6% guinea pigs and 23% chinchillas. There is no evidence of significant differences in the presentation by sex. In rabbits, the diagnosis of the specimens from the group I and II were more common, while in guinea pigs they were II and III. Grade I and II were the most diagnosed in rabbits and guinea pigs. In the radiographs, the most common lesions are the overgrowth of the maxillary and mandibular

roots, elongation of the incisors, loss of the occlusal surface and deformation of the mandibular roots. A method of co-rrection for LL radiographic projections with evident rotation is proposed to maintain the objectivity of the radiographic evaluation

<https://rebiunoda.pro.baratznet.cloud:28443/OpacDiscovery/public/catalog/detail/b2FpOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzQyMzkzODA>

---

**Título:** Prevalencia y hallazgos radiográficos en conejos, cuyos y chinchillas diagnosticados con maloclusión en el Hospital Veterinario de Especialidades en Fauna Silvestre y Etología Clínica de la UNAM, México electronic resource]

**Editorial:** 2020

**Tipo Audiovisual:** maloclusión radiografía *Oryctolagus cuniculus* *Cavia porcellus* Chin-chilla lanigera malocclusion radiography *Oryctolagus cuniculus* *Cavia porcellus* Chinchilla lanigera

**Documento fuente:** Revista de la Facultad de Medicina Veterinaria y de Zootecnia, ISSN 0120-2952, Vol. 67, N°. 1, 2020, pags. 1-16

**Nota general:** application/pdf

**Restricciones de acceso:** Open access content. Open access content star

**Condiciones de uso y reproducción:** LICENCIA DE USO: Los documentos a texto completo incluidos en Dialnet son de acceso libre y propiedad de sus autores y/o editores. Por tanto, cualquier acto de reproducción, distribución, comunicación pública y/o transformación total o parcial requiere el consentimiento expreso y escrito de aquéllos. Cualquier enlace al texto completo de estos documentos deberá hacerse a través de la URL oficial de éstos en Dialnet. Más información: <https://dialnet.unirioja.es/info/derechosOAI> | INTELLECTUAL PROPERTY RIGHTS STATEMENT: Full text documents hosted by Dialnet are protected by copyright and/or related rights. This digital object is accessible without charge, but its use is subject to the licensing conditions set by its authors or editors. Unless expressly stated otherwise in the licensing conditions, you are free to linking, browsing, printing and making a copy for your own personal purposes. All other acts of reproduction and communication to the public are subject to the licensing conditions expressed by editors and authors and require consent from them. Any link to this document should be made using its official URL in Dialnet. More info: <https://dialnet.unirioja.es/info/derechosOAI>

**Lengua:** Spanish

**Enlace a fuente de información:** Revista de la Facultad de Medicina Veterinaria y de Zootecnia, ISSN 0120-2952, Vol. 67, N°. 1, 2020, pags. 1-16

---

## Baratz Innovación Documental

- Gran Vía, 59 28013 Madrid
- (+34) 91 456 03 60
- [informa@baratz.es](mailto:informa@baratz.es)