



A 3D Quantification Technique for Liver Fat Fraction Distribution Analysis Using Dixon Magnetic Resonance Imaging

Instructional and Educational Work

Material Proyactable

This study introduces a unique 3D quantification method for liver fat fraction (LFF) distribution using Dixon Magnetic Resonance Imaging (Dixon MRI). LFF maps, derived from in-phase and water-phase images, are integrated with 3D liver contours to differentiate LFF patterns between normal and steatotic livers, enabling precise assessment of liver fat content.

<https://rebiunoda.pro.baratznet.cloud:28443/OpacDiscovery/public/catalog/detail/b2FpOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzY1MDA5NTA>

Título: A 3D Quantification Technique for Liver Fat Fraction Distribution Analysis Using Dixon Magnetic Resonance Imaging

Editorial: Cambridge JoVE 2023

Descripción física: 1 online resource (1 video file ()) sound, color

Duración: 000000

Mención de serie: Medicine 1940-087X

Intérpretes: Presented by MyJoVE Corporation

Lengua: Presented in English; subtitles in English

Entidades: MyJoVE Corporation production company publisher

Punto acceso adicional serie-Título: Medicine. 1940-087X

Baratz Innovación Documental

- Gran Vía, 59 28013 Madrid
- (+34) 91 456 03 60
- informa@baratz.es

