

Actividad del PH de la solución de ácido orgánico en la obtención de chipas del látex del caucho (Hevea brasiliensis) en la parroquia Patricia Pilar-EcuadorPDFHTML [

2018

text (article)

Analítica

The present research (pH ACTIVITY OF ORGANIC ACID SOLUTION IN OBTAINING SPARKS RUBBER LATEX (Heveabrasiliensis) PATRICIA PILAR COUNTY) was performed taking into account the considerable consumption of formic acid coagulation of rubber latex, which is a pollutant for human beings and the environment, replacing the formic acid with citric acid extracted from lemons that are available to producers and heveicultores. This paper proposed a completely randomized design in factorial arrangement AxB (2x2) with 4 replicates per treatment being the factor A: a1, a2 pH 2.1 y; pH2.6 and factor B: b1; 8% y b2; 12 %. In this research we want to know the influence of pH and the percentage of citric acid in the coagulation of rubber latex through rheological analysis as plasticity, plasticity retention index and viscosity. To evaluate the study factors and their respective levels a multiple range test was executed Tukey 5%. Noting that the treatment with a higher content of plasticity, PRI (Plasticity Retention Index) and viscosity is treatment 2 in the pH level, a percentage of 2.1 using 12% solution. The resulting plasticity of 41.45 and 68.58% of PRI and 100.65 of viscosity. This research was conducted in "Finca Tres Hermanos" located in the 6 ½ km via "El Moral" in the Patricia Pilar County. The experimental responses of the rheological analyzes were performed on the Erco Continental through the Agicom Company, who are dedicated to promoting the cultivation of rubber

The present research (pH ACTIVITY OF ORGANIC ACID SOLUTION IN OBTAINING SPARKS RUBBER LATEX (Heveabrasiliensis) PATRICIA PILAR COUNTY) was performed taking into account the considerable consumption of formic acid coagulation of rubber latex, which is a pollutant for human beings and the environment, replacing the formic acid with citric acid extracted from lemons that are available to producers and heveicultores. This paper proposed a completely randomized design in factorial arrangement AxB (2x2) with 4 replicates per treatment being the factor A: a1, a2 pH 2.1 y; pH2.6 and factor B: b1; 8% y b2; 12 %. In this research we want to know the influence of pH and the percentage of citric acid in the coagulation of rubber latex through rheological analysis as plasticity, plasticity retention index and viscosity. To evaluate the study factors and their respective levels a multiple range test was executed Tukey 5%. Noting that the treatment with a higher content of plasticity, PRI (Plasticity Retention Index) and viscosity is treatment 2 in the pH level, a percentage of 2.1 using 12% solution. The resulting plasticity of 41.45 and 68.58% of PRI and

100.65 of viscosity. This research was conducted in "Finca Tres Hermanos" located in the 6 ½ km via "El Moral" in the Patricia Pilar County. The experimental responses of the rheological analyzes were performed on the Erco Continental through the Agicom Company, who are dedicated to promoting the cultivation of rubber

https://rebiunoda.pro.baratznet.cloud:38443/OpacDiscovery/public/catalog/detail/b2FpOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzY4ODkxODYhcmF0aW9uOmVzLmJhcmF0ei5yZW4vMzY4ODkxODYhcmF0aW9uOmVzLmJhcmF0ei5yZW4vMzY4ODkxODYhcmF0aW9uOmVzLmJhcmF0ei5yZW4vMzY4ODkxODYhcmF0aW9uOmVzLmJhcmF0ei5yZW4vMzY4ODkxODYhcmF0aW9uOmVzLmJhcmF0ei5yZW4vMzY4ODkxODYhcmF0aW9uOmVzLmJhcmF0ei5yZW4vMzY4ODkxODYhcmF0aW9uOmVzLmJhcmF0ei5yZW4vMzY4ODkxODYhcmF0aW9uOmVzLmJhcmF0ei5yZW4vMzY4ODkxODYhcmF0aW9uOmVzLmJhcmF0ei5yZW4vMzY4ODkxODYhcmF0aW9uOmVzLmJhcmF0ei5yZW4vMzY4ODkxODYhcmF0aW9uOmVzLmJhcmF0ei5yZW4vMzY4ODkxODYhcmF0ei5yZW4vMzY4ODkxODyhcmF0ei5yZW4vMzY4ODkxODyhcmF0ei5yZW4vMzY4ODkxODyhcmF0ei5yZW4vMzY4ODkxODyhcmF0ei5yZW4vMzY4ODkxODyhcmF0ei5yZW4vMzY4ODkxODYhcmF0ei5yZW4vMzY4ODkxODYhcmF0ei5yZW4vMzY4ODkxODYhcmF0ei5yZW4vMzY4ODkxODYhcmF0ei5yZW4vMzY4ODkxODYhcmF0ei5yZW4vMzY4ODkxODYhcmF0ei5yZW4vMzY4ODkxODYhcmF0ei5yZW4vMzY4ODkxODYhcmF0ei5yZW4vMzY4ODkxODYhcmF0ei5yZW4vMzY4ODkxODYhcmF0ei5yZW4vMzY4ODkxODYhcmF0ei5yZW4vMzY4ODkxODYhcmF0ei5y

Título: Actividad del PH de la solución de ácido orgánico en la obtención de chipas del látex del caucho (Hevea brasiliensis) en la parroquia Patricia Pilar-EcuadorPDFHTML electronic resource].]

Editorial: 2018

Tipo Audiovisual: Látex Chipas Reológicos Caucho hevea Latex Chipas Rheological Rubber hevea

Documento fuente: Revista Caribeña de Ciencias Sociales (RCCS), ISSN 2254-7630, N°. 7, 2018

Nota general: application/pdf

Restricciones de acceso: Open access content. Open access content star

Condiciones de uso y reproducción: LICENCIA DE USO: Los documentos a texto completo incluidos en Dialnet son de acceso libre y propiedad de sus autores y/o editores. Por tanto, cualquier acto de reproducción, distribución, comunicación pública y/o transformación total o parcial requiere el consentimiento expreso y escrito de aquéllos. Cualquier enlace al texto completo de estos documentos deberá hacerse a través de la URL oficial de éstos en Dialnet. Más información: https://dialnet.unirioja.es/info/derechosOAI | INTELLECTUAL PROPERTY RIGHTS STATEMENT: Full text documents hosted by Dialnet are protected by copyright and/or related rights. This digital object is accessible without charge, but its use is subject to the licensing conditions set by its authors or editors. Unless expressly stated otherwise in the licensing conditions, you are free to linking, browsing, printing and making a copy for your own personal purposes. All other acts of reproduction and communication to the public are subject to the licensing conditions expressed by editors and authors and require consent from them. Any link to this document should be made using its official URL in Dialnet. More info: https://dialnet.unirioja.es/info/derechosOAI

Lengua: Spanish

Enlace a fuente de información: Revista Caribeña de Ciencias Sociales (RCCS), ISSN 2254-7630, N°. 7, 2018

Baratz Innovación Documental

- Gran Vía, 59 28013 Madrid
- (+34) 91 456 03 60
- informa@baratz.es