

## Mini revisión del síndrome de mal ajuste neonatal en equinos

2024

text (article)

Analítica

Neonatal maladjustment syndrome, also called neonatal encephalopathy, perinatal asphyxia, hypoxic-ischemic encephalopathy, and "dumb foal" syndrome, occurs in foals suffering from hypoxia, as well as those suspected of experiencing an altered transition from intrauterine to extrauterine life. However, the etiology is unknown since it may have multiple origins. The primary symptoms include seizures, blindness, vocalizations, wandering, tonic-clonic contractions and loss of the sucking reflex, as well as described in other cases a failure in thermoregulation, seizures and alterations of other body organ systems, such as decreased motility of the gastrointestinal tract. Regarding its pathophysiology, it derives from two moments, the first due to a process of acute cerebral ischemia and energy and oxygen deficit; the second is due to a process of the metabolic neurotransmitter or endocrine imbalances where the inhibitory effects of neurosteroids (adenosine, allopregnanolone, pregnenolone and prostaglandin D2) still present after birth and are believed to be partially responsible for the alteration of consciousness. Treatments usually focus on clinic neonate signs, so pregnancy control in mares and calving accompaniment are essential to prevent this syndrome

Neonatal maladjustment syndrome, also called neonatal encephalopathy, perinatal asphyxia, hypoxic-ischemic encephalopathy, and "dumb foal" syndrome, occurs in foals suffering from hypoxia, as well as those suspected of experiencing an altered transition from intrauterine to extrauterine life. However, the etiology is unknown since it may have multiple origins. The primary symptoms include seizures, blindness, vocalizations, wandering, tonic-clonic contractions and loss of the sucking reflex, as well as described in other cases a failure in thermoregulation, seizures and alterations of other body organ systems, such as decreased motility of the gastrointestinal tract. Regarding its pathophysiology, it derives from two moments, the first due to a process of acute cerebral ischemia and energy and oxygen deficit; the second is due to a process of the metabolic neurotransmitter or endocrine imbalances where the inhibitory effects of neurosteroids (adenosine, allopregnanolone, pregnenolone and prostaglandin D2) still present after birth and are believed to be partially responsible for the alteration of consciousness. Treatments usually focus on clinic neonate signs, so pregnancy control in mares and calving accompaniment are essential to prevent this syndrome

Neonatal maladjustment syndrome, also called neonatal encephalopathy, perinatal asphyxia, hypoxic-ischemic encephalopathy, and "dumb foal" syndrome, occurs in foals suffering from hypoxia, as well as those suspected of experiencing an altered transition from intrauterine to extrauterine life. However, the etiology is unknown since it may have multiple origins. The primary symptoms include seizures, blindness, vocalizations, wandering, tonic-clonic contractions and loss of the sucking reflex, as well as described in other cases a failure in thermoregulation, seizures and alterations of other body organ systems, such as decreased motility of the gastrointestinal tract. Regarding its pathophysiology, it derives from two moments, the first due to a process of acute cerebral ischemia and energy and oxygen deficit; the second is due to a process of the metabolic

neurotransmitter or endocrine imbalances where the inhibitory effects of neurosteroids (adenosine, allopregnanolone, pregnenolone and prostaglandin D2) still present after birth and are believed to be partially responsible for the alteration of consciousness. Treatments usually focus on clinic neonate signs, so pregnancy control in mares and calving accompaniment are essential to prevent this syndrome

https://rebiunoda.pro.baratznet.cloud:38443/OpacDiscovery/public/catalog/detail/b2FpOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzMTA5MDU

**Título:** Mini revisión del síndrome de mal ajuste neonatal en equinos electronic resource].]

Editorial: 2024

**Tipo Audiovisual:** asphyxia encephalopathy neonates foals asfixia encefalopatía neonatos potros asfixia encefalopatía neonatos potros

Documento fuente: Revista Sistemas de Producción Agroecológicos, ISSN 2248-4817, Vol. 15, N°. 2, 2024

(Ejemplar dedicado a: July-December)

Nota general: application/pdf

Restricciones de acceso: Open access content. Open access content star

Condiciones de uso y reproducción: LICENCIA DE USO: Los documentos a texto completo incluidos en Dialnet son de acceso libre y propiedad de sus autores y/o editores. Por tanto, cualquier acto de reproducción, distribución, comunicación pública y/o transformación total o parcial requiere el consentimiento expreso y escrito de aquéllos. Cualquier enlace al texto completo de estos documentos deberá hacerse a través de la URL oficial de éstos en Dialnet. Más información: https://dialnet.unirioja.es/info/derechosOAI | INTELLECTUAL PROPERTY RIGHTS STATEMENT: Full text documents hosted by Dialnet are protected by copyright and/or related rights. This digital object is accessible without charge, but its use is subject to the licensing conditions set by its authors or editors. Unless expressly stated otherwise in the licensing conditions, you are free to linking, browsing, printing and making a copy for your own personal purposes. All other acts of reproduction and communication to the public are subject to the licensing conditions expressed by editors and authors and require consent from them. Any link to this document should be made using its official URL in Dialnet. More info: https://dialnet.unirioja.es/info/derechosOAI

Lengua: Spanish

**Enlace a fuente de información:** Revista Sistemas de Producción Agroecológicos, ISSN 2248-4817, Vol. 15, N°. 2, 2024 (Ejemplar dedicado a: July-December)

## **Baratz Innovación Documental**

- Gran Vía, 59 28013 Madrid
- (+34) 91 456 03 60
- informa@baratz.es