

Presencia de insectos en caraota Phaseolus vulgaris L., sector Mundo Nuevo, parroquia Juan Ángel Bravo, municipio Ezequiel Zamora, Cojedes Venezuela [

2024

text (article)

Analítica

Grain legumes, such as beans Phaseolus vulgaris L., are essential in human nutrition due to their high protein content, complex carbohydrates and low-fat content, being an economical and nutritionally adequate option compared to products of animal origin. In Venezuela, beans are a strategic crop due to their protein content and their ability to adapt to various soils and adverse climates, in addition to improving soil fertility through nitrogen fixation. A study was carried out in the Mundo Nuevo sector, Ezequiel Zamora municipality, Cojedes, whose objective was to identify various pests that affect the bean crop. The results indicate the presence of borer bugs Andrector arcuatus (Olivier) and Andrector ruficornis (Olivier), leafhoppers Empoasca kraemeri (Ross y Moore) and Agrosoma spp.) and thrips (Thrips sp.). These insects cause significant damage, such as leaf perforations, sap sucking, and transmission of viral diseases, negatively affecting crop yields. In addition, beneficial insects were identified such as Podisus sp. (Hemiptera: Pentatomidae), Cycloneda sanguinea (L.) (Coleoptera: Coccinellidae), Diptera: Tachinidae, and Zelus sp. (Hemiptera: Reduviidae). These bioregulators play an important role in the agricultural ecosystem by controlling pest insect populations. Therefore, it is crucial to implement agronomic practices that favor the conservation of these beneficial insects, such as the use of colored traps and natural repellents, instead of chemicals. Beans are not only an essential nutritional source but also a vital crop for food security in Venezuela. However, the presence of various pests represents a considerable challenge for farmers. The identification and proper management of these pests, along with the conservation of beneficial insects, are essential to ensure the sustainability and success of bean cultivation in the Mundo Nuevo sector and other similar regions

Grain legumes, such as beans Phaseolus vulgaris L., are essential in human nutrition due to their high protein content, complex carbohydrates and low-fat content, being an economical and nutritionally adequate option compared to products of animal origin. In Venezuela, beans are a strategic crop due to their protein content and their ability to adapt to various soils and adverse climates, in addition to improving soil fertility through nitrogen fixation. A study was carried out in the Mundo Nuevo sector, Ezequiel Zamora municipality, Cojedes, whose objective was to identify various pests that affect the bean crop. The results indicate the presence of borer bugs Andrector arcuatus (Olivier) and Andrector ruficornis (Olivier), leafhoppers Empoasca kraemeri (Ross y Moore) and Agrosoma spp.) and thrips (Thrips sp.). These insects cause significant damage, such as

leaf perforations, sap sucking, and transmission of viral diseases, negatively affecting crop yields. In addition, beneficial insects were identified such as Podisus sp. (Hemiptera: Pentatomidae), Cycloneda sanguinea (L.) (Coleoptera: Coccinellidae), Diptera: Tachinidae, and Zelus sp. (Hemiptera: Reduviidae). These bioregulators play an important role in the agricultural ecosystem by controlling pest insect populations. Therefore, it is crucial to implement agronomic practices that favor the conservation of these beneficial insects, such as the use of colored traps and natural repellents, instead of chemicals. Beans are not only an essential nutritional source but also a vital crop for food security in Venezuela. However, the presence of various pests represents a considerable challenge for farmers. The identification and proper management of these pests, along with the conservation of beneficial insects, are essential to ensure the sustainability and success of bean cultivation in the Mundo Nuevo sector and other similar regions

https://rebiunoda.pro.baratznet.cloud: 28443/Opac Discovery/public/catalog/detail/b2FpOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPOmNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0aW9uOmVzLmJhcmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0ei5yZW4vMzYzNTg5NjkPomNlbGVicmF0ei5yZW4vMzyZNTg5NjkPomNlbGVicmF0ei5yZW4vMzyZNTg5NjkPomNlbGVicmF0ei5yZW4vMzyZNTg5NjkPomNlbGVicmF0ei5yZW4vMzyZNTg5NjkPomNlbGVicmF0ei5yZW4vMzyZNTg5NjkPomNlbGVicmF0ei5yZW4vMzyZNTg5NjkPomNlbGVicmF0ei5yZW4vMzyZNTg5NjkPomNlbGVicmF0ei5yZW4vMzyZNTg5NjkPomNlbGVicmF0ei5yZW4vMzyZNTg5NjkPomNlbGVicmF0ei5yZW4vMzyZNTg5NjkPomNlbGVicmF0ei5yZW4vMzyZNTg5NjkPomNlbGVicmF0ei5yZW4vMzyZNTg5NjkPomNlbGVicMF0ei5yZW4vMzyZNTg5NjkPomNlbGVicMF0ei5yZW4vMzyZNTg5NjkPomNlbGVicMF0ei5yZW4vMzyZNTg5NjkPomNlbGVicMF0ei5yZW4vMzyZNTg5NjkPomNlbGVicMF0ei5yZW4vMz

Título: Presencia de insectos en caraota Phaseolus vulgaris L., sector Mundo Nuevo, parroquia Juan Ángel Bravo, municipio Ezequiel Zamora, Cojedes Venezuela electronic resource].]

Editorial: 2024

Tipo Audiovisual: insecto plaga caraota identificación incidencia pest insect bean identification incidence

Documento fuente: Ciencia y Tecnología Agropecuaria, ISSN 1900-0863, Vol. 9, N°. 2, 2024, pags. 44-50

Nota general: application/pdf

Restricciones de acceso: Open access content. Open access content star

Condiciones de uso y reproducción: LICENCIA DE USO: Los documentos a texto completo incluidos en Dialnet son de acceso libre y propiedad de sus autores y/o editores. Por tanto, cualquier acto de reproducción, distribución, comunicación pública y/o transformación total o parcial requiere el consentimiento expreso y escrito de aquéllos. Cualquier enlace al texto completo de estos documentos deberá hacerse a través de la URL oficial de éstos en Dialnet. Más información: https://dialnet.unirioja.es/info/derechosOAI | INTELLECTUAL PROPERTY RIGHTS STATEMENT: Full text documents hosted by Dialnet are protected by copyright and/or related rights. This digital object is accessible without charge, but its use is subject to the licensing conditions set by its authors or editors. Unless expressly stated otherwise in the licensing conditions, you are free to linking, browsing, printing and making a copy for your own personal purposes. All other acts of reproduction and communication to the public are subject to the licensing conditions expressed by editors and authors and require consent from them. Any link to this document should be made using its official URL in Dialnet. More info: https://dialnet.unirioja.es/info/derechosOAI

Lengua: Spanish

Enlace a fuente de información: Ciencia y Tecnología Agropecuaria, ISSN 1900-0863, Vol. 9, Nº. 2, 2024, pags. 44-50

Baratz Innovación Documental

- Gran Vía, 59 28013 Madrid
- (+34) 91 456 03 60
- informa@baratz.es